APPLICATION NO. ATTY, DOCKET NO. 09/724,425 10412-026 REFERENCES CITED BY APPLICANT APPLICANT (Use several sheets if necessary) Reed, John C GROUP FILING DATE 1635 11/28/00 U.S. PATENT DOCUMENTS FILING DATE SUBCLASS CLASS NAME IF APPROPRIATE DATE DOCUMENT NUMBER *EXAMINER INITIAL Goodchild 2/21/89 mms 4,806,463 Lee 3/12/91 4,999,290 AB Tsujimoto et al. 05/14/91 AC 5,015,568 Gewirtz et al. 03/24/92 5,098,890 AD Croce 9/22/92 5,149,628 AE Tsujimoto et al. 04/13/93 5,202,429 AF Tsujimoto et al. 10/17/95 5,459,251 AG Hoke et al. 12/17/96 5,585,479 AH3/31/98 Reed JC 5,734,033 Αi Reed JC 6/5/95 5,831,066 Reed JC 5/18/98 6,040,181 AK FOREIGN PATENT DOCUMENTS TRANSLATION SUBCLASS CLASS COUNTRY DATE DOCUMENT NUMBER mas **WIPO** 12/8/94 WO 94/27426 AL PCT 3/30/95 MMS WO 95/08350 **AM** OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.) Asseline U et al., Nucleic acid-binding molecules with high affinity and base sequence specificity: intercalating agents covalently linked to oligodeoxynucleotides. MMS Proc Natl Acad Sci 1984 Jun;81(11):3297-301. Aviv H, Leder P. Purification of biologically active globin messenger RNA by chromatography on AO oligothymidylic acid-cellulose. Proc Natl Acad Sci. 1972 Jun;69(6):1408-12. Bishop JM. The molecular genetics of cancer. Science. 1987 Jan 16;235(4786):305-11. AP Blake KR, Murakami A, Miller PS. Inhibition of rabbit globin mRNA translation by sequence-specific AQ oligodeoxyribonucleotides. Biochemistry. 1985 Oct 22;24(22):6132-8. Boutorin AS et al., Complementary addressed reagents carrying EDTA-FE(II) groups for directed cleavage AR of single-stranded nucleic acids. FEBS Lett. 1984 June;172(1):4346. Chen CH, Sigman DS. Nuclease activity of 1,10-phenanthroline-copper: sequence-specific targeting. AS Proc Natl Acad Sci. 1986 Oct;83(19):7147-51. Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem. 1987 Apr;162(1):156-9.

M Schmelt 6/27/01

Chu BC, Orgel LE. Nonenzymatic sequence-specific cleavage of single-stranded DNA. Proc Natl Acad Sci

ΑT

ΑU

1985 Feb;82(4):963-7.

	AV	Cotter FE, Johnson P, Hall P, Pocock C, al Mahdi N, Cowell JK, Morgan G. Antisense oligonucleotides
nas	~~	suppress B-cell lymphoma growth in a SCID-hu mouse model. Oncogene. 1994 Oct;9(10):3049-55.
	AW	Croce et al., Molecular Basis Of Human B and T Cell Neoplasia. Advance in Viral Oncology, 7:35-51, G. Klein (ed.), New York: Raven Press, 1987.
+	ΔΥ	Haldar S, Beatty C, Tsujimoto Y, Croce CM. The bcl-2 gene encodes a novel G protein. Nature. 1989 Nov
+		9;342(6246):195-8. Holt JT et al., An oligomer complementary to c-myc mRNA inhibits proliferation of HL-60 promyelocytic cells
	AZ	and induces differentiation. Mol Cell Biol. 1988 Feb;8(2):963-73. Kern JA et al., Dexamethasone inhibition of interleukin 1 beta production by human monocytes.
	1 1	Posttranscriptional mechanisms. J Clin Invest. 1988 Jan;81(1):237-44. Kitada S et al., Investigations of antisense oligonucleotides targeted against bcl-2 RNAs. Antisense Research and Development, vol. No. 3, pp. Nos. 157-169, Summer 1993.
	ВВ	Kitada S et al., Reversal of chemoresistance of lymphoma cells by antisense-mediated reduction of bcl-2 gene expression. Antisense Research and Development, vol. No. 4, pp. Nos. 71-79, Summer 1994.
	ВС	Knorre DG et al., Reactive oligonucleotide derivatives and sequence-specific modification of nucleic acids. Biochimie, 1985 Jul-Aug;67(7-8):785-9.
	BD	Matagen (masking tape for gene expression). Anticancer brug box.
	BE	Miyashita T et al., bcl-2 gene transfer increases relative resistance of S49.1 and WEHI7.2 lymphoid cells to cell death and DNA fragmentation induced by glucocorticoids and multiple chemotherapeutic drugs. Cancer Res. 1992 Oct 1;52(19):5407-11.
	BF	Miyashita T et al., Bcl-2 oncoprotein blocks chemotherapy-induced apoptosis in a human leukemia cell line Blood, 1993 Jan 1;81(1):151-7.
	BG	Morvan F et al., alpha-DNA. I. Synthesis, characterization by high field 1H-NMR, and base-pairing properties of the unnatural hexadeoxyribonucleotide alpha-[d(CpCpTpTpCpC)] with its complement beta-[d(GpGpApApGpG)]. Nucleic Acids Res. 1986 Jun 25;14(12):5019-35.
	ВН	Negrini M et al., Molecular analysis of mbcl-2: structure and expression of the murine gene homologous to
+	BI	Paoletti C. Anti-sense oligonucleotides as potential antitumour agents: prospective views and preliminary
\	ВЈ	Reed JC et al. Regulation of chemoresistance by the bcl-2 oncoprotein in non-Hodgkin's lymphoma and
	ВК	Reed et al. Antisense-mediated inhibition of BCL2 protooncogene expression and leukemic cell growth an survival: comparisons of phosphodiester and phosphorothioate oligodeoxynucleotides. Cancer Res. 1990 Oct 15:50(20):6565-70.
	Ві	Reed et al. Suppression of interleukin 2 receptor acquisition by monoclonal antibodies recognizing the 50 KD protein associated with the sheep erythrocyte receptor on human T lymphocytes. J Immunol. 1985 Mar: 134(3):1631-9.
	ВМ	Reed et al. Effect of wheat germ agglutinin on the interleukin pathway of human T lymphocyte activation. J. Immunol. 1985 Jan; 134(1):314-23.
	В	Reed et al. Recombinant interleukin 2 regulates levels of c-myc mRNA in a cloned murine T lymphocyte. Mol Cell Biol. 1985 Dec;5(12):3361-8.

M Schwelt 6/27/01

mas	во	Reed et al. BCL2-mediated tumorigenicity of a human T-lymphoid cell line: Synergy with MYC and inhibition by BCL2 antisense. Proc Natl. Acad. Sci., 87:3660-3664 (1990).
1		Reed JC et al., Regulation of bcl-2 proto-oncogene expression during normal human lymphocyte proliferation. Science. 1987 Jun 5;236(4806):1295-9.
	BQ	Stein CA, Cohen JS. Oligodeoxynucleotides as inhibitors of gene expression: a review. Cancer Res. 1988 May 15;48(10):2659-68.
	BR	Stein CA et al., Physicochemical properties of phosphorothioate oligodeoxynucleotides. Nucleic Acids Res
	BS	Strasser A et al., bcl-2 transgene inhibits T cell death and perturbs thymic self-censorship. Cell. 1991 Nov 29;67(5):889-99.
	вт	Stull RA et al., Antigene, ribozyme and aptamer nucleic acid drugs: progress and prospects. Pharm Res. 1995 Apr;12(4):465-83.
	BU	Tsujimoto Y et al., Analysis of the structure, transcripts, and protein products of bcl-2, the gene involved in human follicular lymphoma. Proc Natl Acad Sci, 1986 Jul;83(14):5214-8.
	BV	Uhlmann, Eugen et al."Antisense Oligonucleotides: A New Therapeutic Principle," Chemical Reviews, vol. 90, No. 4, p. Nos. 544-584, Jun. 1990.
	BW	van den Elsen P et al., Isolation of cDNA clones encoding the 20K T3 glycoprotein of human T-cell receptor complex. Nature. 1984 Nov 29-Dec 5;312(5993):413-8.
	вх	Vlassov VV et al., Complementary addressed modification and cleavage of a single stranded DNA fragme with alkylating oligonucleotide derivatives. Nucleic Acids Res. 1986 May 27;14(10):4065-76.
	BY	Webb TR et al., Hybridization triggered cross-linking of deoxyoligonucleotides. Nucleic Acids Res. 1986 Oct 10;14(19):7661-74.
	BZ	gene. J Immunol. 1987 Apr 1;138(7):2169-76.
	CA	Wickstrom EL et al., Human promyelocytic leukemia HL-60 cell proliferation and c-myc protein expression are inhibited by an antisense pentadecadeoxynucleotide targeted against c-myc mRNA. Proc Natl Acad St. U.S.A. 1988 Feb;85(4):1028-32.
	СВ	Williams GT. Programmed cell death: apoptosis and oncogenesis. Cell. 1991 Jun 28;65(7):1097-8.
	cc	1986 Jun;83(12):4143-6.
V	CE	Zamecnik PC et al., Inhibition of Rous sarcoma virus replication and cell transformation by a specific
EXAMINE	ĒR ,	M Schwolf DATE CONSIDERED 6/27/01

NY2 - 1166583.1